

ENUMERATIVE GEOMETRY OF DIVISORIAL FAMILIES OF RATIONAL CURVES

ZIV RAN

May 5, 2002/ Rev. June 10, 2002

ABSTRACT. We compute the number of irreducible rational curves of given degree with 1 tacnode or 1 triple point in $\mathbb{P}^2$ or 1 node in $\mathbb{P}^3$ meeting an appropriate generic collection of points and lines. As a byproduct, we also compute the number of rational plane curves of degree d passing through $3d - 2$ given points and tangent to a given line. The method is 'classical', free of Quantum Cohomology.

In the past 15 years or so a number of classical problems in the enumerative geometry of curves in $\mathbb{P}^n$ were solved, first for $n = 2$, any genus [R1], then for any n , genus 0. The latter development was initiated by Kontsevich-Manin who developed and used the rather substantial machinery of Quantum Cohomology (cf. e.g. [FP]). Subsequently, in a series of papers [R2-R5] the author developed an elementary alternative approach, free of Quantum cohomology, and used it to solve a number of classical enumerative problems for rational, and sometimes elliptic, curves in $\mathbb{P}^n$, $n \geq 2$. The present paper continues this series. The object here is to enumerate the irreducible rational curves of given degree d in $\mathbb{P}^2$ with one tacnode or one triple point passing through $3d - 2$ general points (see Theorems 2,5 below), as well as the irreducible rational curves in $\mathbb{P}^3$ with one ordinary node which contain a general points and are incident to $4d - 2a - 1$ general lines (see Thm 1 below). Note that the family of 1-tacnodal (resp. 1-triple point, resp. 1-nodal) curves in $\mathbb{P}^2$ (resp. $\mathbb{P}^2, \mathbb{P}^3$) is of codimension 1 in the family of all rational curves so that we are effectively computing the 'degree', in a sense, of certain natural divisors in the family of all rational curves. Indeed by a result of Diaz and Harris [DH] in the case of $\mathbb{P}^2$, the general member

Updates and corrections available at math.ucr.edu/~ziv/papers/1nodal.pdf.
Research Partially supported by NSA Grant MDA904-02-1-0094; reproduction and distribution by US government permitted.

of any such divisor, if not nodal, is either 1-cuspidal (which case was enumerated in [R2]) or 1-tacnodal or has 1 triple point. It seems very likely, but doesn't seem to be in the literature, that the natural analogue of this result also holds for any $n \geq 3$: i.e. that the general member of any divisor in the family of rational curves is either smooth, or 1-nodal reducible (any n), or 1-nodal irreducible ($n=3$). As a byproduct of the proof of Theorem 2, we also obtain a formula for one of the 'characteristic numbers' for rational plane curves, viz. the number of such curves of degree d passing through $3d - 2$ general points and tangent to a given line (cf. Cor. 4 below).

Enumeration of rational plane curves with a tacnode or triple point appears to be well within the range of interest, at least, of classical geometers. For quartics, results of this kind, at least the enumeration of quartics with a triple point, were obtained by Zeuthen in 1882. Recently, enumerative results of this kind for curves with 'few' singularities on general surfaces were obtained by Kleiman and Piene [KP]. Enumeration of 1-nodal rational curves in $\mathbb{P}^n$ for any n was recently announced by Zinger [Z], using Quantum Cohomology. As we shall see below, the 1-nodal case in $\mathbb{P}^3$ case is analogous to, but easier than the 1-tacnodal and 1 triple-point cases in $\mathbb{P}^2$. Our proof is based on the intersection calculus on the nonsingular model of the surface swept out by the appropriate 1-parameter family of rational curves, developed in earlier papers [R2-R4], together with elementary residual-intersection (in particular, double-point) theory as in [F]

We begin by reviewing some qualitative results about families of rational curves in $\mathbb{P}^n$, especially for $n = 2$ or 3 . See [R2][R3] [R4] and references therein for details and proofs. In what follows we denote by $\bar{V}_d$ the closure in the Chow variety of the locus of irreducible nonsingular rational curves of degree d in $\mathbb{P}^n, n = 3$, with the scheme structure as closure, i.e. the reduced structure (recall that the Chow form of a reduced 1-cycle Z is just the hypersurface in $G(1, \mathbb{P}^3)$ consisting of all linear spaces meeting Z); if $n = 2$ $\bar{V}_d$ is just the (closure of) the Severi variety. Thus $\bar{V}_d$ is irreducible reduced of dimension

$$\begin{aligned} \dim(\bar{V}_d) &= 4d, n = 3 \\ \dim(\bar{V}_d) &= 3d - 1, n = 2. \end{aligned}$$

Let

$$A_1, \dots, A_k \subset \mathbb{P}^n$$

be a generic collection of linear subspaces of respective codimensions $a_1, \dots, a_k$, $2 \leq a_i \leq n$ (so if $n = 2$ these are just points). We denote by

$$B = B_d = B_d(a.) = B_d(A.)$$

the normalization of the locus (with reduced structure)

$$\{(C, P_1, \dots, P_k) : C \in \bar{V}_d, P_i \in C \cap A_i, i = 1, \dots, k\}$$

which is also the normalization of its projection to $\bar{V}_d$, i.e. the locus of degree- d rational curves (and their specializations) meeting $A_1, \dots, A_k$. We have

$$(1) \quad \dim B = (n+1)d + n - 3 - \sum (a_i - 1).$$

When $\dim B = 0$ we set

$$(2) \quad N_d(a.) = \deg(B).$$

When $n = 2$, so all the $a_i = 2$ they will be dropped. The integer k is called the *length* of the condition-vector $(a.)$. The numbers N_d and $N_d(a.)$, first computed in general by Kontsevich and Manin (see for instance [FP] and references therein), are computed in [R2],[R3] by an elementary method based on recursion on d and k .

Now suppose $\dim B = 1$ and let

$$(3) \quad \pi : X \rightarrow B$$

be the normalization of the tautological family of rational curves, and

$$f : X \rightarrow \mathbb{P}^n$$

the natural map. The following summarizes results from [R2][R3][R4] :

Theorem 0. (i) X is smooth .

(ii) Each fibre C of π is either

(a) a $\mathbb{P}^1$ on which f is either an immersion with at most one exception which maps to a cusp ($n = 2$) or an embedding ($n > 2$); or

(b) a pair of $\mathbb{P}^1$'s meeting transversely once, on which f is an immersion with nodal image ($n = 2$) or an embedding ($n > 2$); or

(c) if $n = 3$, a $\mathbb{P}^1$ on which f is a degree-1 immersion such that $f(\mathbb{P}^1)$ has a unique singular point which is an ordinary node.

(iii) If $n > 2$ then $\bar{V}_{d,n}$ is smooth along the image $\bar{B}$ of B , and $\bar{B}$ is smooth except, in case some $a_i = 2$, for ordinary nodes corresponding to curves meeting some A_i of codimension 2 twice. If $n = 2$ then $\bar{V}_{d,n}$ is smooth in codimension 1 except for a cusp along the cuspidal locus and normal crossings along the reducible locus, and $\bar{B}$ has the singularities induced from $\bar{V}_{d,n}$ plus ordinary nodes corresponding to curves with a node at some A_i , and no other singularities.

Next, we review some of the enumerative apparatus introduced in [R3][R4] to study X/B . Set

$$(4) \quad m_i = m_i(a.) = -s_i^2, i = 1, \dots, k.$$

Note that if $a_i = a_j$ then $m_i = m_j$; in particular for $n = 2$ they are all equal and will be denoted by m_d . It is shown in [R2] [R3][R4] that these numbers can all be computed recursively in terms of data of lower degree d and lower length k . For instance for $n = 2$ we have

$$(5) \quad 2m_d = \sum_{d_1+d_2=d} N_{d_1} N_{d_2} d_1 d_2 \binom{3d-4}{3d_1-2}.$$

For $n > 2$, note that

$$s_i \cdot s_j = N_d(\dots, a_i + a_j, \dots, \hat{a}_j, \dots), i \neq j.$$

(so for $n = 2$ this is always 0). Also, letting R_i denote the sum of all fibre components not meeting s_i , we have

$$(6) \quad s_1 \cdot R_2 = \sum N_{d_1}(A^1, A_1, \mathbb{P}^{s_1}) N_{d_2}(A^2, A_2, \mathbb{P}^{s_2}).$$

the summations being over all $d_1 + d_2 = d$, $s_1 + s_2 = 3$ and all decompositions $A = (A_1, A_2) \coprod (A^1) \coprod (A^2)$ (as unordered sequences or partitions); similarly for the other $s_i \cdot R_j$. So all these numbers may be considered known. Then we have

$$m_i = \frac{1}{2}(s_i \cdot R_j + s_i \cdot R_p - s_j \cdot R_p) - s_i \cdot s_j - s_i \cdot s_p + s_j \cdot s_p$$

for any distinct i, j, p , and the RHS here is an expression of lower degree and/or length, hence may be considered known.

Next, set

$$L = f^*(\mathcal{O}(1)),$$

and note that

$$L^2 = N_d(2, a.), \quad L \cdot s_i = N_d(a_1, \dots, a_{i+1}, \dots), \quad i = 1, \dots, k$$

(in particular, $L \cdot s_i = 0$ if $a_i = n$.) We computed in [R3] that, for any i ,

$$L \sim ds_i - \sum_{F \in \mathcal{F}_i} \deg(F) F + (N_d(a_1, \dots, a_i + 1, \dots) + dm_i) F_0$$

where F_0 is the class of a complete fibre and $\mathcal{F}_i$ is the set of fibre components not meeting s_i . Consequently we have

$$(7) \quad N_d(2, a_1, \dots) = 2dN_d(a_1 + 1, a_2, \dots) + d^2 m_1(a.) - \sum_{F \in \mathcal{F}_1(a.)} (\deg F)^2$$

and clearly the RHS is a lower degree/length expression, so all the $N_d(2, \dots)$ are known. We also have for $n > 2$ that

$$N_d(a_1, a_2 + 1, \dots) - N_d(a_1 + 1, a_2, \dots) =$$

$$(8) \quad dN_d(a_1 + a_2, \dots) - \sum_{F \in (\mathcal{F}_1 - \mathcal{F}_2)(a.)} (\deg F) + N_d(a_1 + 1, a_2, \dots) + dm_1(a.)$$

and again the RHS here is 'known', hence so is the LHS, which allows us to 'shift weight' between the a_i 's till one of them becomes 2, so we may apply (7), and thus compute all of the $N_d(a.)$'s.

Next, it is easy to see as in [R3] that

$$(9) \quad L.R_i = \sum_{d_1 + d_2 = d} \binom{3d-1}{3d_1-1} d_1 d_2^2 N_{d_1} N_{d_2}, \quad n = 2$$

(in this case this is independent of i and we will just write it as $L.R$);

$$(10) \quad L.R_i = \sum d_2 N_{d_1}(a.^1, s_1) N_{d_2}(a.^2, s_2), \quad n > 2$$

the summation for $n > 2$ being over all $d_1 + d_2 = d, s_1 + s_2 = 3$ and all decompositions

$$A. = (A.^1) \coprod (A.^2)$$

(as unordered sequences or partitions) such that $A_i \in (A.^1)$.

Finally, the relative canonical class $K_{X/B} = K_X - \pi^*(K_B)$ was computed in [R3] as

$$(11) \quad K_{X/B} = -2s_i - m_i F + R_i$$

for any i . Note that $-R_i^2$ equals the number of reducible fibres in the family X/B , a number we denote by $N_d^{\text{red}}(a.)$, and which is easily computable by recursion. From this we compute easily that

$$(12) \quad L.K_{X/B} = -2N_d(\dots a_i + 1 \dots) - dm_i + L.R_i,$$

$$(13) \quad K_{X/B}^2 = -N_d^{\text{red}}(a.).$$

Now in case $n = 3$, for any condition-vector $(a.)$ of weight

$$\sum (a_i - 1) = 4d - 1,$$

we denote by $N_d^\times(a.)$ the number of 1-nodal irreducible rational curves in P^3 meeting a generic collection of linear spaces of respective codimensions $(a_1, \dots, a_k)$.

Theorem 1. *We have for any $i = 1, \dots, k$,*

$$(14) \quad N_d^\times(a.) =$$

$$(d-3)N_d(2, a.) - N_d(\dots a_i + 1 \dots) - (4d+2)m_i + 2L.R_i + N_d^{\text{red}}(a.).$$

proof. We set things up so as to apply the *double point formula* (in a relative form [F]). Let

$$X_B^2$$

denote the fibre square of X/B , and

$$\Delta \simeq X \subset X_B^2$$

the diagonal. Then X_B^2 is smooth except at points (p, p) , where p is a singular point of a fibre of X/B . Locally at such points, if X/B is given locally at p by

$$xy = t,$$

then X_B^2 is given by

$$x_1 y_1 = x_2 y_2 = t,$$

and therefore has an ordinary 3-fold double point. Moreover the diagonal Δ is defined by

$$x_1 = x_2, y_1 = y_2$$

and in particular is a non-Cartier divisor. Let

$$(15) \quad b : Y \rightarrow X_B^2$$

denote the blow-up of Δ , with exceptional divisor

$$\Delta' = b^*(\Delta).$$

Then it is easy to see that b is a small resolution of X_B^2 with exceptional locus

$$E = \sum E_p$$

consisting of a $\mathbb{P}^1$ for each relatively singular point p . Moreover the fibre of Y over B corresponding to a reducible fibre $C_1 \cup_p C_2$ of X/B is of a 'honeycomb' shape

$$\begin{array}{ccc} B_{(p,p)}C_1^2 & \cup & C_1 \times C_2 \\ C_2 \times C_1 & \cup & B_{(p,p)}C_2^2 \end{array}$$

where

$$B_{(p,p)}C_1^2 \cap B_{(p,p)}C_2^2 = E_p, C_1 \times C_2 \cap C_2 \times C_1 = \emptyset.$$

Also $\Delta' \rightarrow \Delta$ is just the blowing up of all the points p . Furthermore, the identity of rational functions, locally at p ,

$$\frac{y_2 - y_1}{x_2 - x_1} = \frac{y_1}{x_2} = \frac{y_2}{x_1}$$

shows that there $b : Y \rightarrow X_B^2$ coincides with the blowup of the ideal (y_1, x_2) , and with that of the ideal (y_2, x_1) .

We will need to know the normal bundle

$$\nu = N_{\Delta'/Y}.$$

To this end, note that, clearly

$$\omega_{\Delta'} = b^* \omega_X(E).$$

On the other hand, the map (15) is small, hence crepant, so

$$(16) \quad \omega_Y = b^* \omega_{X_B^2} = b_1^* \omega_X \otimes b_2^* \omega_X \otimes c^* \omega_B^{-1}$$

where we use the evident maps

$$b_i : Y \rightarrow X, i = 1, 2, c : Y \rightarrow B.$$

Therefore by the adjunction formula we conclude

$$(17) \quad \nu = b^* \omega_{X/B}^{-1}(E).$$

Now we are ready to apply the double-point formula to the map

$$f^2 = (f_1, f_2) : X_B^2 \rightarrow \mathbb{P}^3 \times \mathbb{P}^3.$$

This shows that the cycle of ordered pairs (x_1, x_2) in X/B^2 such that $f(x_1) = f(x_2)$ is residual to Δ' in $f^{2*}(\Delta_{\mathbb{P}^3})$. Therefore, as in ([CR], Thm 3) we find that

$$N_d^\times(a.) = \frac{1}{2} c_3(f_1^*(\mathcal{O}(1)) \otimes f_2^*(Q) \otimes \mathcal{O}(-\Delta')),$$

where Q denotes the universal quotient bundle on $\mathbb{P}^3$. Then a straightforward calculation, based on the intersection calculus reviewed above, yields the formula (14). $\square$

We turn next to the enumeration of rational plane curves with a tacnode. Denote by N_d^{tac} (resp. κ_d) the number of rational curves with a tacnode (resp. cusp) passing through $3d - 2$ generic points in $\mathbb{P}^2$, and recall that κ_d was computed in [R3].

Theorem 2. *We have*

$$(18) \quad N_d^{\text{tac}} = (d-4)N_d - (d-7)(d^2 + 7d - 4)m + \frac{d-7}{2}L.R + 2N_d^{\text{red}} + \frac{1}{2}\kappa_d$$

proof. We use an analogous setup and notation as above, this time for a pencil of rational curves in $\mathbb{P}^2$ through $3d-2$ points. To begin with, we describe an algebraic setup for the (relative) Gauss mapping. Consider the relative principal parts sheaf for L on X/B , which fits in an exact sequence

$$(19) \quad 0 \rightarrow \Omega_{X/B}(L) \rightarrow P_{X/B}(L) \rightarrow L \rightarrow 0.$$

Setting $V = H^0(\mathcal{O}_{\mathbb{P}^2}(1))$, we have a natural map $V \rightarrow P_{X/B}(L)$ which combines with the Euler sequence to yield an exact diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & f^*\Omega_{\mathbb{P}^2}(1) & \rightarrow & V \otimes \mathcal{O}_X & \rightarrow & f^*\mathcal{O}_{\mathbb{P}^2}(1) & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \parallel & & \\ 0 & \rightarrow & \Omega_{X/B}(L) & \rightarrow & P_{X/B}(L) & \rightarrow & L & \rightarrow & 0. \end{array}$$

Let P' denote the pushout of $P_{X/B}(L)$ by the natural map

$$\Omega_{X/B}(L) \rightarrow \omega_{X/B}(L).$$

Now let $b : X'/B \rightarrow X/B$ denote the blowup of all singular points of fibres and all 'cuspidal' points, i.e. all points $(x, b) \in X$ such that $f(x)$ is a cusp on $f(X_b)$ (see [R3] for a discussion and enumeration of these). Let $E = \sum E_p$ be the exceptional divisor over the singular points of fibres and $U = \sum U_q$ be the exceptional divisor over the cuspidal points. Set $f' = b \circ f$. Then it is easy to see by a local computation that

Lemma 3. *The image of the natural map*

$$f'^*\Omega_{\mathbb{P}^2} \rightarrow b^*\omega_{X/B}$$

coincides with $b^\omega_{X/B}(-E - U)$*

proof. It suffices to prove locally at each fibre node or cuspidal point p that

$$\text{im}(f^*\Omega_{\mathbb{P}^2} \rightarrow \omega_{X/B}) = \omega_{X/B}.I_p.$$

In local coordinates at p , the family X/B is given by

$$xy = t$$

with $f = (x, y)$ and $\omega_{X/B}$ is generated by $dx \wedge dy \cdot dt^{-1}$. Since

$$dx \wedge dt = xdx \wedge dy, dy \wedge dt = -ydx \wedge dy,$$

we have that $f^*\Omega_{\mathbb{P}^2}$ is generated by $xdx \wedge dy \cdot dt^{-1}, ydx \wedge dy \cdot dt^{-1}$, as claimed.

In the cuspidal case, our family has local coordinates u, t with

$$f = (u^2, (u^2 - t)u) = (x, y).$$

Here $\omega_{X/B}$ is generated by du and its subsheaf generated by

$$dx = 2udu, dy = (3u^2 - t)du$$

clearly coincides with the subsheaf generated by udu and tdu , as claimed $\square$

It follows from Lemma 3 that if we let P denote the image of the natural map

$$V \otimes \mathcal{O}_{X'} \rightarrow b^*P',$$

then we get a diagram with exact rows and columns

$$\begin{array}{ccccccc} 0 & \rightarrow & f'^*\Omega_{\mathbb{P}^2}(1) & \rightarrow & V \otimes \mathcal{O}_X & \rightarrow & f^*\mathcal{O}_{\mathbb{P}^2}(1) \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ (19) \quad 0 & \rightarrow & b^*\omega_{X/B}(L - E - U) & \rightarrow & P & \rightarrow & b^*L \rightarrow 0. \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

Now let

$$\phi : I \rightarrow \mathbb{P}^2$$

be the incidence (or flag) variety, with universal flag

$$V \otimes \mathcal{O}_I = F^0 \supset F^1 = \phi^*\Omega_{\mathbb{P}^2}(1) \supset F^2 \supset (0),$$

$$F^0/F^1 = \phi^*\mathcal{O}(1).$$

Now the diagram (19) gives rise to a lifting of f' to a morphism

$$g : X' \rightarrow I$$

with

$$(20) \quad g^*(F^0/F^1) = L' := b^*(L), g^*(F^1/F^2) = b^*(\omega_{X/B})(-E - U).$$

Clearly, the value of g at a point

$$(x, b) \in X' \setminus E \setminus U$$

mapping to

$$y \in \mathbb{P}^2$$

is the pair

$$(y, T_y(f(X_b)))$$

consisting of y and the tangent line to $f(X_b)$ at $f(y)$; in particular, the tacnodes in the family X/B correspond (1:2) to the double points of g , which we propose to count as in the proof of Theorem 1. To this end, the construction of a 'good' desingularization of $X' \times_B X'$ must be modified to take into account the fact that each E_p appears in its fibre over B with multiplicity 2.

Consider then the fibre square

$$\bar{Y} = X' \times_B X'.$$

Its singular points are as follows. First, for each cuspidal point q on X , we get a 3-fold ODP (y, y) where y is the intersection of U_q with the proper transform of the fibre of X/B containing q . We desingularize this by blowing up $U_q \times U_q$ (which is the same locally as blowing up the diagonal). Here the construction is exactly as in the proof of Thm 1.

Additionally, $\bar{Y}$ is singular along each $E_p \times E_p$, where it has a local equation of the form

$$x_1^2 u_1 = x_2^2 u_2$$

where $x = 0$ is the equation of E_p , u is a coordinate on E and x_i, u_i are the pullbacks of x, u . This can be desingularized by blowing up the locus

$$x_1 = x_2 = 0,$$

i.e. $E_p \times E_p$. The exceptional divisor maps to $E_p \times E_p$ generically with degree 2, ramified over the 4 rulings

$$E_p \times \{0, \infty\} \cup \{0, \infty\} \times E_p,$$

where $0, \infty$ are the 2 intersections of E_p with the other 2 components of its fibre, and having $\mathbb{P}^1$ fibres over $\{0, \infty\}^2$.

Let

$$b : Y \rightarrow \bar{Y}$$

be the global desingularization thus obtained, and $\Delta'' \subset Y$ the proper (=total) transform of the diagonal. It is easy to see that the map

$$b : \Delta'' \rightarrow X'$$

identifies Δ'' with the blowup X'' of all singular points of set-theoretic fibres of X/B (i.e. all the points of the form (y, y) or $(0, 0)$ or (∞, ∞) in the above notation.

The normal bundle

$$\nu = N_{\Delta''/Y}$$

can be computed much like before: first,

$$\omega_{\Delta''} = b^* \omega_{X'}(E'')$$

where $E'' = \sum E''_r$ is the exceptional divisor of $b|_{\Delta''}$. Next, as before,

$$\omega_{\bar{Y}} = b_1^* \omega_X \otimes b_2^* \omega_X \otimes \omega_B^{-1}.$$

As $\bar{Y}$ has an ordinary double surface generically along $\bigcup_p E_p$, it follows that

$$\omega_Y = b^*(\omega_{\bar{Y}})(-\bigcup_p E_p).$$

Putting these together we see that, using divisor notation and setting $K = K_X - K_B$,

$$(21) \quad \nu = -(b^*(\omega_{X'} - \omega_B - E) - E'') = -(K + U - E'').$$

Consequently, we have

$$(22) \quad \nu^2 = K^2 - 2N_d^{\text{red}} - 2\kappa_d$$

where κ_d is the number of cuspidal rational plane curves of degree d through $3d - 2$ generic points, computed in [R3]. Of course, we also have

$$(23) \quad L.\nu = -L.K = 2dm - L.R$$

Now, to use double-point theory consider the cartesian product $I^2 = I \times I$ with projections

$$p_i : I^2 \rightarrow I, i = 1, 2, \phi^2 : I^2 \rightarrow (\mathbb{P}^2)^2.$$

Let $F_\cdot$ be the dual filtration to $F^\cdot$, defined by

$$F_i = (F^0/F^i)^* \subset (F^0)^*.$$

In I^2 , $\phi^{2*} \Delta_{\mathbb{P}^2}$ is defined as the zero-locus of a map

$$p_1^* F_1 \rightarrow p_2^*(F_3/F_1),$$

and inside this, Δ_I is defined as the zero- locus of a map

$$p_1^*(F_2/F_1) \rightarrow p_2^*(F_3/F_2).$$

Considering as before the map

$$f^2 = (f_1, f_2) : Y \rightarrow I^2,$$

double-point (or more precisely, residual-intersection) theory shows that

$$2N_d^{\text{tac}} =$$

$$c_2(f_1^*(F_1^*) \otimes f_2^*((V^* \otimes \mathcal{O})/F_1) \otimes \mathcal{O}(-\Delta'')) c_1(f_1^*((F_2/F_1)^*) \otimes f_2^*(F_3/F_2) \otimes \mathcal{O}(-\Delta'')).$$

Using (20), the c_2 factor can be identified, writing $L_i = p_i^* L$, $K_i = p_i^*(\omega_{X/B})$, as

$$L_2^2 + L_1 L_2 - L_2 \Delta' + (L_1 - \Delta')^2$$

while the c_1 factor is

$$(K_1 - E - U) + L_2 - \Delta''.$$

Then a routine computation, using as above the necessary intersection theory on Y and X yields the formula (18). $\square$

As a bonus for the construction of the lift g above- and without using any double-point theory, we get a formula for one of the 'characteristic numbers', namely the number N_d^t of rational plane curves through $3d - 2$ points tangent to a given line. Thus let

$$p_2 : I \rightarrow \mathbb{P}^{2v}$$

be the natural map of the incidence variety to the dual projective plane, and set

$$f^v = p_2 \circ g : X' \rightarrow \mathbb{P}^{2v}, L^v = f^{v*} \mathcal{O}(1).$$

As above, we can write

$$L^v \sim (2d - 2)s_1 - \sum_{F \in \mathcal{F}_1} (\deg(F) - 1)F' - \sum E_p - \sum U_q + xF_0$$

where the first sum is over all fibre components of X/B not meeting s_1 , F' denotes the proper transform of F on X' , the second and third sums cover over all exceptional divisors of X'/X , F_0 denotes a fibre of X'/B , and we

have simply used the fact that the dual to a rational curve of degree d has degree $2d - 2$. Now intersect this expression with s_1 and note that

$$L^v.s_1 = N_{d\rightarrow},$$

i.e. the number of degree- d rational curves through $A_1, \dots, A_{3d-2}$ with a given tangent direction at A_1 . This yields

$$x = N_{d\rightarrow} + (2d - 2)m_d.$$

Since $N_d^t = \mathcal{L}^{v^2}$, this number can now be easily computed from our intersection calculus, yielding the following

Corollary 3. *We have*

$$N_d^t = 4(d-1)^2 m_d + d(d-1)N_{d\rightarrow} + N_d^{\text{red}} - \kappa_d - 2 \sum_{F \cap s_1 = \emptyset} (\deg(F) - 1)^2.$$

Finally, we take up the problem of enumerating the triple points, i.e. of computing the number N_d^{tri} of irreducible rational curves with a triple point passing through $3d-2$ generic points. Again our tool will be a suitable residual-intersection computation. Using notations as above, take 3 copies

$$Y_{12}, Y_{13}, Y_{23}$$

of the small resolution of X_B^2 constructed above. We have evident 'first projection' maps

$$Y_{12} \rightarrow X, Y_{13} \rightarrow X$$

which allows us to construct

$$Y_{12} \times_X Y_{13}$$

together with a map

$$p_{23} : Y_{12} \times_X Y_{13} \rightarrow X_B^2.$$

Let Z be the unique component of $Y_{12} \times_X Y_{13}$ which dominates X_B^2 . Define

$$Y^*$$

as the blowup of Z in the proper transform of the diagonal in X_B^2 . Then Y^* comes equipped with projections

$$p_{ij} : Y^* \rightarrow Y_{ij},$$

a map

$$f^3 = (f_1, f_2, f_3) : Y^* \rightarrow (P^2)^3$$

as well as with a birational map

$$b^* : Y^* \rightarrow X_B^3.$$

An elementary analysis shows that Y^* has exactly 2 singular points, each isomorphic to a cone over a Segre variety $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$, hence also to the germ at the origin of the determinantal variety $M_{2 \times 3}^1$ of (2×3) matrices of rank at most 1, and consequently each of these points admits a small 'determinantal' resolution with $\mathbb{P}^1$ as exceptional locus. It can be shown that this resolution admits a natural morphism to the relative Hilbert scheme $\mathcal{Hilb}_3(X/B)$. Though we could replace Y^* by this resolution, this will not be necessary since by definition the big diagonals are already Cartier on Y^* .

Now consider on Y^* the locus

$$D_{12} = p_{12}^* D$$

where $D \subset Y$ is the double locus of f^2 , i.e. the closure of the locus of distinct point (y_1, y_2) in the same fibre which map to a node, i.e. such that $f(y_1) = f(y_2)$. Then clearly

$$[D_{12}] = p_{12}^*([D]) = c_2(Q_2(L_1 - \Delta_{12}))$$

where

$$Q_i = f_i^* Q, L_i = f_i^* L, \Delta_{12} = p_{12}^*(\Delta').$$

Now consider the intersection $D_{12}.D_{13}$. This consists of the locus we want, i.e. that of ordered distinct triples (y_1, y_2, y_3) on the same fibre such that $f(y_1) = f(y_2) = f(y_3)$, plus the locus $D_{12}.\Delta_{23}$. Then residual-intersection theory tells us that

$$6N_d^{\text{tri}} = c_2(Q_2(L_1 - \Delta_{12})).c_2(Q_3(L_1 - \Delta_{13} - \Delta_{23})).$$

This expression can be computed by routine calculations, using the intersection calculus developed above; the only possibly nonobvious terms are

$$\Delta_{12}^2 \Delta_{13}^2 = \Delta_{12}^2 \Delta_{23}^2 = K^2,$$

$$\Delta_{12}^2 \Delta_{13} \Delta_{23} = (K - E)^2.$$

The former follows from the fact that

$$p_{12*}(\Delta_{13}^2) = -p_{12*}(p_{13}^*(K_1 - E)) = -K_1,$$

and that $\Delta_{12}^2 = -(K_1 - E)$ (cf. (17)). The latter follows from the fact that

$$p_{12*}(\Delta_{13} \Delta_{23}) = \Delta_{12}.$$

This yields

Theorem 5. *We have*

$$6N_d^{\text{tri}} = (3d^2 - 18d + 30)N_d + (3d - 18)(dm_d - L.R) - 6N_d^{\text{red}}$$

Check. For $d = 4$, we have $m_d = 428$, $N_d = 620$, $L.R = 3276$, $N_d^{\text{red}} = 2124$, therefore

$$N_4^{\text{tri}} = 60,$$

a number computed classically by Zeuthen and recently, with modern methods, by Kleiman and Piene [KP]. I am grateful to Steve Kleiman for this information, which led to a correction of an error in an earlier statement of the formula.

REFERENCES

- [CR] M.C.Chang and Z. Ran, *Dimension of families of space curves*, Comp. Math. **90** (1994), 53-57.
- [DH] S. Diaz and J. Harris, *Geometry of Severi varieties*, Trans. Amer. Math. Soc. **309** (1988), 1-34.
- [F] W. Fulton, *Intersection theory*, Springer, 1984.
- [FP] ——— and R. Pandharipande, *Notes on stable maps and quantum cohomology*, Algebraic Geometry (Santa Cruz 1995).
- [KP] S. Kleiman, R. Piene, *Enumerating singular curves on surfaces*, Algebraic geometry, Hirzebruch 70, Contemporary Math., vol. 241, 1999, pp. 209-238, corrections and revision in math.AG/9903192.
- [R1] Z.Ran, *Enumerative geometry of singular plane curves*, Invent. math **97** (1987), 447-465.
- [R2] ———, *Bend, break and count*, Isr. J. Math **111** (1999), 109-124.
- [R3] ———, *Bend, break and count II*, Math. Proc. Camb. Phil. Soc. **127** (1999), 7-12.
- [R4] ———, *On the variety of rational space curve*, Isr. J. Math **122** (2001), 359-370.
- [R5] ———, *The degree of the divisor of jumping rational curves*, Quart. J. Math. (2001), 1-18.
- [Z] A. Zinger, *Enumeration of 1-nodal rational curves in projective spaces*, math.AG **0204236**.

UNIVERSITY OF CALIFORNIA, RIVERSIDE
E-mail address: `ziv@math.ucr.edu`